## Pain Management Unit QEII Health Sciences Centre Opioid risk flow sheet

Admission date:																				
Risk factors  □ personal history addiction □ smoker □ family history addiction □ criminal history □ history of psychiatric illness □ pain from mva □ preadolescent sexual abuse □ lack of family supports □ age < 45 years							Opioid attitudes  ☐ insists on specific opioid ☐ insists on IR opioids ☐ not interested in non-opioid therapy ☐ intolerant to non-opioid pharmacotherapy ☐ reports that only opioids work													
Aberrant Drug-Related Behaviours	admission																			
requests early refills																				
increased dose without authorization														1					1	
reports lost or stolen prescription														l						
changes route of administration: IV/nasal																				
hoarding opioid																				
non-adherent to non-opioid therapy																				
attempts to get opioid from other doctors																				
getting opioid from non-medical sources																				
forging prescriptions																				
uses opioid for sedation																				
uses opioid for euphoria																				
using opioid in response to stressors																				
abuses alcohol or illicit drugs																				
drug related crime																				
involved with street drug culture																				
selling or giving away opioid																				
opioid withdrawal symptoms																				
involved in car or other accident																				
multiple contacts between appointments																				
missed appointments																				
Observations																				
withdrawal																				
intoxication	+																			
sedation																				
unkempt																				
concerns from significant others	+-																			
	+-																			